Mazuri® Small Bird Diet

(Available at www.mazuri.com or through a Mazuri® retailer.)



Formula Code - 56A6

Description

Mazuri® Small Bird Diet is a diet for small psittacines and passerines with diverse particle sizes. This diet is specially formulated for non-breeding birds (young birds after first molt and up to breeding age, and for birds kept in non-breeding situations). This diet is not for growing chicks.

Features and Benefits

- Mixed particle size Supports natural foraging behavior.
- **Better than seeds** Less waste, no sorting, consistent nutrition.
- Contains ground flaxseed A source of essential omega-3 fatty acids to support overall health.

| No added artificial colors or artificial flavors. Contains antioxidant vitamin E – An essential nutrient. | | 经总额 |
|--|--------------------|-----|
| Product Form Extruded feed: 2 particle sizes. | Catalog # | 为证法 |
| 2.5 lb. resealable poly bag 25 lb. net weight paper sack | 0054885 0001452 | 经商务 |

Guaranteed Analysis

| Crude protein not less than 14.5% | Ash not more than 8.0% | | |
|---|---|--|--|
| Crude fat not less than 5.0% | Vitamin E not less than150 IU/lb | | |
| Crude fiber not more than 5.0% | Omega-3 fatty acids not less than 0.50% | | |
| Moisture not more than 12.0% | | | |
| Total Microorganisms, min | 270,000 CFU/g | | |
| (Lactobacillus acidophilous, Lactobacillus casei, Bifidobacterium thermophilum, Enterococcus faecium) | | | |
| Contains a source of live (viable) naturally occurring microorganisms. | | | |

Ingredients

Ground Corn, Wheat Middlings, Dehulled Soybean Meal, Corn Gluten Meal, Ground Flaxseed, Glyceryl Monostearate, Dicalcium Phosphate, Soybean Oil, Calcium Carbonate, Wheat Germ. Brewers Dried Yeast, Calcium Propionate (a preservative), Salt, L-Lysine, DL-Methionine, Dried Lactobacillus acidophilus Fermentation Product, L-Ascorbyl-2-Polyphosphate (Vitamin C), Choline Chloride, Dried Lactobacillus casei Fermentation Product, Dried Bifidobacterium thermophilum Fermentation Product, Pyridoxine Hydrochloride, Biotin, Dried Enterococcus faecium Fermentation Product, d-Alpha Tocopheryl Acetate (Vitamin E), Menadione Sodium Bisulfite Complex (Vitamin K), Cholecalciferol (Vitamin D3), Preserved with Mixed Tocopherols, Rosemary Extract, Folic Acid, Vitamin A Acetate, Citric Acid (a Preservative), Riboflavin Supplement, Manganous Oxide, Zinc Oxide, Calcium Iodate, Calcium Pantothenate, Thiamine Mononitrate, Beta Carotene, Nicotinic Acid, Vitamin B12 Supplement, Copper Sulfate, L-Tryptophan, Zinc Sulfate, Sodium Selenite, Cobalt Carbonate.

Feeding Directions

Feed free choice to non-breeding birds and provide fresh food daily.

- A 3 $\frac{1}{2}$ oz (100 g) bird will typically eat between 2 to 3 Tbsp (10 15 g) per day, while a 1 lb. (450 g) bird may consume between 1/3 to 1/2 cup (33 – 40 g) per day.
- Fresh fruits, vegetables and seeds may be provided, but no more than 20% of the total diet.
- To switch birds to this diet, gradually replace your old diet with this diet at the rate of 10% per day. Do not overfeed during this transition time and monitor your bird to be sure it is eating the new diet.
- Thoroughly wash feed and water bowls on a regular basis. It is always good practice to wash hands thoroughly after feeding and/or handling pets. This diet is not for human consumption.

12/23/2020 56A6-RCHM-W13

Mazuri® Small Bird Maintenance Diet

Approximate Nutrient Composition¹

MIITDIENTS

| NUTRIENTS | |
|------------------------|------|
| Protein, % | 15.0 |
| Arginine, % | 0.79 |
| Cystine, % | |
| Glycine, % | 0.61 |
| Histidine, % | 0.38 |
| Isoleucine, % | 0.64 |
| Leucine, % | |
| Lysine, % | 0.75 |
| Methionine, % | 0.48 |
| Phenylalanine, % | 0.75 |
| Threonine, % | |
| Tryptophan, % | 0.15 |
| Tyrosine, % | 0.47 |
| Valine, % | 0.70 |
| | |
| Fat (Ether extract), % | |
| Linoleic acid, % | |
| Linolenic acid, % | |
| Omega-3 Fatty Acids, % | |
| Omega-6 Fatty Acids, % | 2.24 |
| | |
| Fiber (Crude), % | |
| NDF, % | 15 |
| ADF, % | 4 |
| | |
| Ash, % | 5.0 |

| WACKU WINEKALS | |
|---------------------------------|------|
| Calcium, % | 0.90 |
| Chloride, % | 0.24 |
| Magnesium, % | 0.19 |
| Phosphorus, % | 0.70 |
| Phosphorus (non-phytate), % | |
| Potassium, % | |
| Sodium, % | |
| Sulfur, % | |
| | |
| TRACE MINERALS | 40 |
| Copper, ppm | 12 |
| lodine, ppm | 1.34 |
| Iron, ppm | |
| Manganese, ppm | |
| Selenium (added), ppm | |
| Zinc, ppm | 99 |
| VITAMINS | |
| Ascorbic acid, ppm | 280 |
| Biotin, ppm | |
| Choline, ppm | |
| Folic acid, ppm | |
| Niacin, ppm | |
| Pantothenic acid, ppm | |
| Pyridoxine, ppm | |
| Riboflavin, ppm | |
| Thiamin, ppm | |
| Vitamin B ₁₂ , µg/kg | |
| | |
| Vitamin A, IU/kg | |
| Vitamin D ₃ , IU/kg | |
| Vitamin E, IU/kg | 176 |
| Vitamin K (as menadione), ppm | 2.9 |
| Beta-Carotene, ppm | 9.6 |

Storage Conditions

For best results, ensure polybag seal is fully closed during storage, or store contents of open paper sack in container with sealing lid. Store in a cool (75°F or colder), dry (approximately 50% RH) location free from rodents and insects. Do not offer moldy or insect-infested feed to animals as it may result in illness, performance loss or death. Freezing will not harm the diet and may extend freshness. Adhere to the "Best if used by" date or use within 1 year of bag manufacturing.

Mazuri® is a registered trademark of PMI Nutrition International, LLC.

12/23/2020 56A6-RCHM-W13

¹ Based on the latest ingredient analysis information. Since nutrient composition of ingredients varies, analyses will vary accordingly.